

CALADIUM ROOT ROT CAUSED BY PYTHIUM MYRIOTYLUM

W. H. Ridings and R. D. Hartman¹

Caladiums (*Caladium hortulanum* Birdsey), well known for their attractive multi-colored foliage, are primarily used for indoor pot plants and home landscaping (8). Native to the Amazon Basin, caladiums are a 2-million-dollar industry in Florida, which produces approximately 95% of the world's commercial crop (3,8). There are over 2000 named varieties with approximately 50 varieties produced commercially (8).

Several fungi (i.e., *Sclerotium rolfsii* Sacc., *Rhizoctonia solani* Kuehn, and *Pythium* spp.), either together or separately, have been associated with root rot of caladium (1). Recently, *Pythium myriotylum* Drechs. was shown to cause severe root rot of the caladium cultivar *Candidum* and proved to be the most aggressive species of *Pythium* tested (7),

SYMPTOMS. *P. myriotylum* apparently infects only the roots of caladium resulting in reddish brown lesions, root rot, and severe root loss (fig. 1A, B). With the cultivar *Candidum* corm piece germination is slowed in infested soil, and plant development is restricted with as great as a 41% loss in corm weight (7).




Fig. 1. Caladium roots infected with *P. myriotylum*. A) Healthy roots (left) and diseased roots (right). B) Diseased roots showing reddish brown lesions (l) and root rot (r).

¹Plant Pathologist, Pan-American Plant Co., P.O. Box 428, West Chicago, IL 60185.

Disease Development. Besides being present in soil, *P. myriotylum* has been detected in decayed roots attached to non-hot water treated corms being used for planting (1). A soil temperature of 32 C (90 F) and high soil moisture favor the development of this fungus significantly over a soil temperature of 24 C (75 F) and high moisture (7). Although not tested on caladium, *P. myriotylum* has been shown to act synergistically with other soil-borne pathogens (2).

Host Range. Other than caladium, *P. myriotylum* is known to attack a wide variety of crops including tomato, cucumber, bean, tobacco, watermelon, eggplant, papaya, peanut, rye, and other grasses (5,9).

CONTROL. The general procedures outlined for control of other soil pests will serve to control *P. myriotylum*. These procedures consist of 1) selection of a well-drained planting site, 2) fumigation of soil with a general fumigant such as methyl bromide, 3) hot water treatment of corms before planting (4,6), and 4) prevention of the introduction of contaminated soil into the planting site either from equipment or flooding rains. If *P. myriotylum* is detected despite the above precautions, then a soil drench with a fungicide such as Truban 30 WP may be helpful for control. Varieties resistant to root rot have not been researched.

References Cited

1. Bureau of Plant Pathology (Plant Disease Records), Fla. Dept. Agr. and Consumer Serv., Div. Plant Ind., Gainesville. (Unpublished)
2. Garcia, R., and D. J. Mitchell. 1975. Synergistic interaction of *P. myriotylum* with *Fusarium solani*, and *Meloidogyne arenaria* in pod rot of peanut. *Phytopathology* 65:832-833.
3. Holms, L. L., J. Hendry, L. Tubbs, A. L. Hall, and D. Pittman. 1965. Caladium bulbs. Highlands County DARE Rept. for 1965-1975. 11 p.
4. Knauss, J. F. 1975. Description and control of *Fusarium* tuber rot of caladium. *Plant Dis. Reptr.* 59:975-979.
5. McCarter, S. M., and R. H. Littrell. 1968. Pathogenicity of *Pythium myriotylum* to several grass and vegetable crops. *Plant Dis. Reptr.* 52:179-183.
6. Rhoades, H. L. 1964. Effect of hot water treatment of seed tubers and soil fumigation for control of root knot on yield of caladiums. *Plant Dis. Reptr.* 48:568-571.
7. Ridings, W. H., and R. D. Hartman. 1976. Pathogenicity of *Pythium myriotylum* and other species of *Pythium* to caladium derived from shoot-tip culture *Phytopathology* 66: (In press).
8. Sheehan, T. J. 1967. Caladium production in Florida. Univ. Fla., IFAS, Gainesville. Circ. 128B. 7 p.
9. Waterhouse, G. M., and J. M. Waterston. 1966. *Pythium myriotylum*. CMI Descriptions of Pathogenic Fungi and Bacteria 278. Commonwealth Mycological Institute, Kew, Surrey, England. 2 p.